

Excision/Mohs Wound Care Instructions

Dressing Materials:

- Telfa or non-adherent dressing pads.
- 4 x 4 gauze or cotton pads (like those for removing make-up).
- Micropore paper tape or Band-aids.
- Topical ointment such as Vaseline, petroleum, or Aquaphor.
- Do NOT use polysporin, Neosporin, bacitracin, or other antibiotic ointments unless a prescription antibiotic is prescribed by your provider.

Other: _____

Wound Care:

1. If steri-strips or surgical glue is applied to the surgical site, no wound care is needed. If the steri-strips fall off during the time the sutures are in place use the wound care instructions provided below.
2. Leave the original dressing in place for **24 hours, 48 hours, or 72 hours**; DO NOT get the dressing wet.
3. After _____ hours wash your hands and remove the dressing. If there is crusting or scabbing then get in the shower and soak it off, but DO NOT forcibly rub the crust off the wound.
4. Gently cleanse the wound with one of the following: soap and water, saline, or _____ and pat dry with a clean gauze, cotton pad, or a soft, clean washcloth.
5. Apply a thin layer of ointment (Vaseline or Aquaphor) or prescription antibiotic on all suture lines using a clean, new Q-tip.
6. Cover the wound with a non-stick bandage, such as a Telfa, and secure with paper tape. For smaller areas, a Band-aid may be used, but make sure the sticky part of the Band-aid does not touch the wound directly.
7. Repeat your dressing changes (steps 3-6) once or twice daily depending on the amount of drainage until the sutures are removed or the wound is healed.
8. As long as your drainage is not pus-like (thick, yellow, sometimes smelly), then it is unlikely to be infected.

Activities After Surgery:

- For _____ hours after surgery, DO NOT get the dressing wet.
- For 48 hours after surgery, avoid alcohol consumption.
- For 1 week after surgery, avoid any activity that stretches your wound, increases your heart rate, or rubs against your dressing. Avoid sports, vacuuming, bending below your waist, or lifting more than 10 lbs.
- Face surgery: DO NOT bend your head below your waist. Keep your head elevated with 1 or 2 extra pillows when sleeping or lying down.
- Arm or leg surgery: Raise the limb as much as possible above your waist when you are resting.

- Baths/ swimming / hot tubs: Avoid submerging the wound under water until the sutures have been removed.
- Discontinue smoking. Smoking slows wound healing, increases risk of infection, and may result in a more prominent scar.
- Remember to use broad-spectrum sunscreen with Zinc or Titanium Dioxide and at least a SPF of 30 or higher once the area has healed.

Pain, Swelling, Redness, Bruising, Drainage:

- It is normal to have some swelling, redness, and/or bruising—all of which will improve over time (days to weeks).
- Expect a “black eye” on both sides if any procedure was performed above the nose—this is normal. If there is swelling, please use a cold compress and, if directed, take OTC Benadryl unless allergic or a provider has instructed you to not take Benadryl due to other medical conditions.
- Elevating the wound on extra pillows will help reduce swelling in the affected area.
- Applying a cold compress (bag of frozen peas) will also help to reduce swelling in the affected area.
- Once you get home, hold the ice pack against the dressing for 10 minutes several times per hour. Continue this until bedtime and even the following day. DO NOT apply the cold compress directly on your skin.
There may be some clear wound draining, which should stop after a few days.
- The drainage may be pink or yellow in color, which is also normal.
- The drainage may last longer if the wound was left open to heal and not stitched closed.
- If your pain is not controlled with cold compresses, you may take extra strength Tylenol (acetaminophen 500 mg) as needed unless you are allergic, you have liver disease, or a provider has instructed you to not take acetaminophen due to other medical conditions.
- If your pain is not relieved with cold compresses AND extra strength Tylenol (acetaminophen 500 mg, DO NOT EXCEED 2.5 GRAMS PER DAY), then please call the office. Please see below for contact number during normal business hours and after business hours.

Bleeding After Surgery:

- Bleeding after surgery is uncommon but may be seen in the first 48 hours.
- If you are bleeding, please follow the instructions below:
 1. Lie down, elevate the wound if possible.
 2. Apply continuous, firm pressure for 15 minutes. Do not stop applying pressure until 15 minutes have passed. Repeat up to 3 times.
 3. If the bleeding has soaked your dressing, it should be removed and a new one should be placed.
 4. If the bleeding has not stopped after 3 applications of firm, uninterrupted pressure contact our office. Refer to the contact numbers provided below for the phone number during normal business hours and the after-hours phone number.

When Should I Call the Office?

- Fever of 101° Fahrenheit or 38.3° Celsius or higher.
- Bleeding is not controlled by direct pressure.
- Pain that increases each day or is not relieved by over-the-counter medications.
- Rapid or increasing swelling, heat, and pain around the wound.
- Reopening of the wound at any time.
- Increased or expanding redness and warmth around the wound, or pus draining from the wound.
- There is always an on-call provider after hours. Should you experience any of the above symptoms, call the after-hours phone number listed below and our on-call service will contact a healthcare provider from _____, who can answer your questions and provide guidance. Please call to help avoid a visit to the emergency room.
 - **After office hours, contact (____) _____ - _____ ext. _____.**
- However, if you are unable to reach a provider and you have an urgent issue concerning your surgical site/wound, please call 911 or go to the emergency room.

Should you experience any of the following signs or symptoms as listed above, please contact us at:

- **During office hours, contact (____) _____ - _____ ext. _____.**
- **After office hours, contact (____) _____ - _____ ext. _____.**